arXiv:0907.5225v3 [math.SG] 24 Mar 2010

OPEN GROMOV-WITTEN THEORY ON CALABI-YAU THREE-FOLDS I

VITO IACOVINO

Abstract. We propose a general theory of the Open Gromov-Witten invariant on Calabi-Yau three-folds. We introduce the moduli space of multi-curves and show how to get invariants from it. Our construction is based on an idea of Witten.

In the special case that each connected component of the Lagrangian submanifold has the rational homology of a sphere we define rational numbers $F_{g,h}$ for each genus g and h boundary components.

1. Introduction

Let M be a Calabi-Yau three-fold and let L be a special Lagrangian submanifold of M. The open Gromov-Witten invariants of the pair (M, L) are supposed to count the holomorphic bordered curves of M with boundary mapped into L, or equivalently the Euler Characteristic of the moduli space of pseudo holomorphic bordered curves. Mathematicians have been unable to give a general construction of these invariants because the moduli space of pseudo holomorphic bordered curves has codimension one boundary, making its Euler Characteristic ill defined. A mathematical definition was given only assuming particular symmetry of the pair (M, L) (see [5], [6]).

Witten in [7] defines the Open Topological string partition function for a pair (X, L) that, by analogy with the closed case (it is well known that the closed topological string partition function can be computed in terms of closed Gromov-Witten invariants), it should correspond to the Open Gromov-Witten invariants. These invariants have also been computed in string theory in many examples. In this paper and the following [3] we give a mathematical construction of the invariant of [7]. This leads to a mathematical definition of Open Gromov-Witten invariants.

The boundary of the moduli space of pseudo holomorphic disks is of codimension one because of the bubbling of disks and the bubbling of spheres from a constant disk. The second problem is less harmful since it can happen only in discrete points of the space of parameters (such as the space of compatible almost complex structures) that we need to define the moduli space of disks. To deal with the bubbling off of spheres from a constant disk we assume that L is homologically trivial (we do not assume that L is connected). This condition is also natural in string theory since it implies that the topological charge of the D-brane is zero. It was also used in mathematics independently by Joyce.

The problem of the splitting of the curves is a much more delicate issue since it happens along the entire space of parameters, making meaningless the count of the number of curves. In fact, because of this the number of disks strongly depends

---

Date: version: March 25, 2010.

on the perturbation of the pseudo-holomorphic equation. Our idea to solve this problem is based on a result of [7]. Witten argues that the open topological string partition function receive contributions also from degenerate or partial degenerate curves (that is objects that are made by usual curves joined by infinite thin strips living on L). Witten proposes that this contribution can be computed in terms of Chern-Simons integrals. In the case of the cotangent bundle $M = T^*L$, there are no non-constant holomorphic curves with boundary mapped into the zero section. The Open Topological String is equivalent to the Chern-Simons theory on L (the degenerate curves correspond to Feynman graphs of the Chern-Simons theory). For more general Calabi-Yau (where pseudo-holomorphic curves are present) Witten computes the contribution of the degenerated curves using Wilson loops integrals associated to the boundary of the curves.

From the work of Witten follows that open topological string theory does not simply count open curves, but it counts what we define as multi-curves (in analogy with the physical terminology multi-instantons). Roughly a multi-disk is described by a tree with a disk associated to each vertex. The moduli space of multi-curves provide the natural mathematical setting for the computation of [7]. In this setting the problem of the splitting of the curves is completely solved.

In the particular case that L has the rational homology of a sphere there is a nice interpretation of the invariant in terms of generalized linking numbers. In fact the Wilson loop integrals of [7] can be formulated in terms of linking number of the boundary circles of the curves. This computation is a sum of contributions of multi-disks. The contribution of a multi-disk to the Open Gromov-Witten invariant is computed in terms of the linking numbers of the boundaries components of the disks vertices.

The solution of the problem of the bubbling off of the disks can be explained roughly as follows. In a one parameter family of compatible almost complex structures of M, a disk A can split into two disks B and C. The contribute of the 2-disk B, C is given by the linking number of the boundary knots. When the disk A disappear this linking number of B, C jumps. Therefore the jump of the contribution of the 2-disk compensates the jump of the number of 1-disks (that is ordinary disks). Similarly, disappearing of an n-disk is compensated by the discontinuity in the m-disk contribution for $m > n$.

In [4], we apply the analysis of this paper to non-compact situations. This includes a mathematical definition of the invariants arising in Large N duality.

In the following paper [3] we consider the case that $H_1(L, \mathbb{Q}) \neq 0$. We construct the Open Gromov-Witten potential S as an homotopy class of solutions of the Master equation in the ring of the functions on $H^*(L, \mathbb{Q})$ with coefficients in the Novikov ring (this is the effective action of the Open Topological String of [7]). S is defined up to master homotopy (that is unique up to equivalence). The evaluation of S on its critical points leads to numerical invariants.

The paper is organized as follows. In section two we define the moduli space of multi-disks as a Kuranishi space. It is easy to see that an orientation of the moduli space of disks induces an orientation on the moduli space of multi-disks. We show that the boundary faces can be identified in pairs in a way compatible with the orientation.

In section three we define system of chains. In section four we introduce the notion of coherent perturbation in the boundary. With the constraint of coherence in the boundary the evaluation map in the punctures of multi-disks leads to a system of chain well defined up to homotopy. This procedure is a minor adaptation of the procedure developed in [1].

In section five we specialize to the case that each connected component of L has the rational homology of a sphere. In this case (generalizing the usual definition of linking number of curves) we can assign to our system of chains a well defined rational number. This rational number is the Open Gromov-Witten invariant $F_{0,1}$.

In section six we consider the higher genus case. Most of we have said for multi-disks can be adapted to multi-curves. In the case that each connected component of L has the rational homology of a sphere we define the Open Gromov-Witten invariants $F_{g,h} \in \mathbb{Q}$ of genus g and h boundary components.

Acknowledgements. We are grateful to D. Joyce for many usefull correspondence.

2. Multi-disks

Let M be a Calabi-Yau three-fold and let L be a special Lagrangian submanifold of M (or more generally a Lagrangian with Maslov index zero). In this section we define the moduli space of multi-disks with boundary mapped in L. For this we need to introduce first some notations.

Given a ribbon tree T we denote by $V(T)$ the set of vertices of T, by $E(T)$ the set of edges of T. $H(T)$ is the set of edges of T with an assigned orientation, or equivalently the set of half-edges of T (for each internal edge there correspond two elements of $H(T)$ and for each external edge there is one element of $H(T)$, the one starting in the vertex). For $v \in V(T)$, $H(v)$ denotes the set of oriented edges starting from v.

Definition 1. *A decorated tree is a ribbon tree T with a relative homological class $A_v \in H_2(M, L)$ assigned to each vertex v. If $A_v \neq 0$ the vertex v can have arbitrary valence. If $A_v = 0$ the valence of v has to be at least 3.*

The relative homology class of T is by definition the sum of the homology class of its vertices $A = \sum_{v \in V(T)} A_v$. We denote with $\mathcal{T}$ the set of decorated trees, $\mathcal{T}(A)$ the set of decorated trees in the homology class A and $\mathcal{T}_k$ the set of decorated trees with k external edges.

Let $C_2(L)$ be the compactification of the configuration space of two points of L. Recall that $C_2(L)$ is the manifold with boundary given by the geometric blow-up along the diagonal Δ of $L \times L$. The boundary is a S^2 bundle on Δ

$$\partial C_2(L) \to \Delta \cong L. \tag{1}$$

In fact $C_2(L)$ is isomorphic to the sphere bundle $S(TL)$ of L.

For each internal edge $e \in E(T)$, denote by $C_e(L)$ a copy of $C_2(L)$ where we consider the points identified with the half edges associated to e. An orientation of e induces an identification between $C_e(L)$ and $C_2(L)$. For each external edge $e \in E(T)$, define $C_e(L) = L$. The projection $C_2(L) \to L^2$ induces a map

$$\prod_{e \in E(T)} C_e(L) \to L^{H(T)}. \tag{2}$$

Fix a compatible almost-complex structure J. Let $\overline{\mathcal{M}}_n(A)$ be the moduli space of pseudo-holomorphic disks with n marked points with a fixed cyclic order in the relative homology class $A \in H_2(X, L)$ (this is usual denoted as the main component in [2]). By Chapter 7 of [2], $\overline{\mathcal{M}}_n(A)$ can be endowed with a weakly submersive Kuranishi structure with corners.

For each $v \in V(T)$ denote by $\overline{\mathcal{M}}_v$ a copy of $\overline{\mathcal{M}}_{H(v)}(A_v)$ where we consider the boundary marked points labeled (respecting the cyclic order of the ribbon structure) with the half edges $H(v)$ starting in v.

The evaluation on the punctures defines a strongly continuous map

$$\prod_{v \in V(T)} \overline{\mathcal{M}}_v \to L^{H(T)}. \tag{3}$$

The moduli space of T-multi-disks $\overline{\mathcal{M}}_T$ is given by taking the fiber product of the maps (3) and (2) mod-out by the automorphism group of the decorated tree

$$\overline{\mathcal{M}}_T = \left(\prod_{v \in V(T)} \overline{\mathcal{M}}_v \times_{L^{H(T)}} \prod_{e \in E^(T)} C_e(L) \right) / \mathrm{Aut}(T).$$

Since we assume that the Kuranishi structure of each $\overline{\mathcal{M}}_v$ is weakly submersive, the space $\overline{\mathcal{M}}_T$ has a natural Kuranishi structure (see Section A1.2 of [2]). Moreover this Kuranishi structure has a tangent space.

2.1. **Boundary.** Fix a decorated tree T. For each $v \in V(T)$ the boundary of $\overline{\mathcal{M}}_v$ can be decomposed in different components

$$\partial \overline{\mathcal{M}}_{H(v)}(A_v) = \left(\bigsqcup_{\substack{H_1 \cup H_2 = H(v) \\ A_1 + A_2 = A}} \overline{\mathcal{M}}_{\{H_1 \cup *\}}(A_1) \times_L \overline{\mathcal{M}}_{\{H_2 \cup *\}}(A_2) \right) / \mathbb{Z}_2 \tag{4}$$

corresponding to all the ways to subdivide $H(v)$ in two sets. In the particular case that $H(v)$ is empty there is an extra term coming from sphere bubble attached to constant disks:

$$\partial \overline{\mathcal{M}}_0 = (\overline{\mathcal{M}}_1 \times_L \overline{\mathcal{M}}_1)/\mathbb{Z}_2 \sqcup \overline{\mathcal{M}}_{0,1} \times_M L. \tag{5}$$

Here $\overline{\mathcal{M}}_{0,1}$ is the moduli space of spheres with one marked point. The last term of (5) comes from the bubbling of the spheres from constant disks. It has been discussed in section 32.1 of [2] (these are the boundary nodes of type E in Definition 3.4 of [5]).

For each pair (T, v) (with $v \in V(T)$) define

$$\partial_v \overline{\mathcal{M}}_T = \left(\left(\partial \overline{\mathcal{M}}_v \times \prod_{v' \neq v} \overline{\mathcal{M}}_{v'} \right) \times_{L^{H(T)}} \prod_e C_e(L) \right) / \mathrm{Aut}(T, v) \tag{6}$$

and for each pair (T, e) (with $e \in E(T)$ an internal edge)

$$\partial_e \overline{\mathcal{M}}_T = \left(\left(\prod_v \overline{\mathcal{M}}_v \right) \times_{L^{H(T)}} \left(\partial C_e(L) \times \prod_{e' \neq e} C_{e'}(L) \right) \right) / \mathrm{Aut}(T, e). \tag{7}$$

The boundary of $\overline{\mathcal{M}}_T$ can be subdivided in the Kuranishi spaces $\partial_v \overline{\mathcal{M}}_T$, $\partial_e \overline{\mathcal{M}}_T$.

Observe that $\partial_v \overline{\mathcal{M}}$ is union of boundary faces of $\overline{\mathcal{M}}$. The elements of each boundary face come with a special nodal point. Let

$$\mathrm{ev}_* : \partial_v \overline{\mathcal{M}}_T \to L \tag{8}$$

be the evaluation on the special nodal point. Let $S(\partial_v\overline{\mathcal{M}}_T)$ be the the Kuranishi space given by the pull-back of the sphere bundle (1) using the map (8):

$$S(\partial_v\overline{\mathcal{M}}_T) = (\mathrm{ev}_*)^*(S(TL)).$$

The natural map

$$S(\partial_v\overline{\mathcal{M}}_T) \to \partial_v\overline{\mathcal{M}}_T \tag{9}$$

defines an S^2-fibration of Kuranishi spaces in the sense that locally $S(\partial_v\overline{\mathcal{M}}_T)$ is a product as Kuranishi space of $\partial_v\overline{\mathcal{M}}_T$ and the two dimensional sphere S^2.

Definition 2. *Let $e \in E(T)$ be an internal edge attached to the vertices v_1, v_2. The decorated tree T/e is the tree obtained by contracting e to a vertex. The relative homology class of the vertex e/e is the sum of the homology classes of the vertices v_1 and v_2, $A_{e/e} = A_{v_1} + A_{v_2}$.*

The Kuranishi space $\partial_v\overline{\mathcal{M}}_T$ is subdivided in boundary faces of $\overline{\mathcal{M}}_T$. In fact we have:

Lemma 1. *Let T_0 be the tree with no edges and one vertex v. There is an isomorphism of Kuranishi spaces:*

$$S(\partial_v\overline{\mathcal{M}}_{T_0}) = \partial_e\overline{\mathcal{M}}_{T'} \sqcup S(\overline{\mathcal{M}}_{0,1} \times_M L) \tag{10}$$

where T' is the graph given by two vertices connected by an edge e. If T has at least one edge, there is an isomorphism of Kuranishi spaces:

$$S(\partial_v\overline{\mathcal{M}}_T) \cong \bigsqcup_{(T'/e,e/e)=(T,v)} \partial_e\overline{\mathcal{M}}_{T'} \tag{11}$$

where the union is over all the trees T' and edges $e \in E(T')$ such that $(T'/e, e/e) = (T, v)$.

Proof. The lemma is immediate from the definition. Consider first (10) and let T' be the graph given by two vertices w, z connected by an edge e. We have

$$\partial_e\overline{\mathcal{M}}_{T'} = ((\overline{\mathcal{M}}_w \times \overline{\mathcal{M}}_z) \times_{L^2} \partial C_2(L))/\mathbb{Z}_2$$

and equation (5)

$$\partial_v\mathcal{M}_0 = (\overline{\mathcal{M}}_1 \times_L \overline{\mathcal{M}}_1)/\mathbb{Z}_2 \sqcup \overline{\mathcal{M}}_{0,1} \times_M L.$$

All the fiber products are made using the evaluation map as usual. Observe that

$$(\mathcal{M}_1 \times_L \mathcal{M}_1) = (\mathcal{M}_1 \times \mathcal{M}_1) \times_{L^2} \Delta$$

where Δ is the diagonal of L^2. The formula (10) follows since in this case the map (8) is the natural projection on $\Delta \cong L$. In the same way (11) follows from (4). □

2.2. **Orientation.** Let $T \in \mathcal{T}$ be a decorated tree. For each internal edge $e \in E(T)$, an orientation of e induces an orientation of $C_e(L)$ and an identification $C_e(L) \cong C_2(L)$. Let $o_e \cong \mathbb{Z}_2$ be the set of orientations of the edge e. Let $o_{\mathrm{ex}} \cong \mathbb{Z}_2$ be the set of parities of the ordering of the external edges of T. Define

$$o_T = (\otimes_{e\in E(T)} o_e) \otimes o_{\mathrm{ex}}.$$

Assume that L is relative spin and oriented. For each vertex $v \in V(T)$, let $\mathcal{M}_{0,v}$ be a copy of the moduli space of disks without punctures in the homology class A_v. By Section 44 of [2], $\mathcal{M}_{0,v}$ has a natural orientation.

We can identify $\mathcal{M}_T$ with an open subset of

$$\prod_{v\in V(T)} \mathcal{M}_{0,v} \times \prod_{e\in E(T)} \partial D_e$$

where ∂D_e is the boundary of the disk from where the edge e starts. From this the following Lemma follows:

Lemma 2. *An orientation of the moduli space of disks induces an isomorphism between the orientation bundle of $\mathcal{M}_T$ and o_T.*

With this orientation we can prove that the identifications of the boundary faces given in Lemma 1 are compatible with the orientation.

Lemma 3. *Endow $\partial_e\mathcal{M}_T$ with the orientation induced as boundary face of $\mathcal{M}_T$ and endow $S(\partial_v\overline{\mathcal{M}}_T)$ with the orientation induced by the fibration (9). Then the isomorphisms (11) and (10) reverse the orientation.*

Proof. This is analogous to the proof of Proposition 46 of [2]. We use the same convention of Section 45 of [2]. We need to explain the lemma only in the particular case that the graph T is given by two vertices v, w connected by an edge e. The tangent space on $\mathcal{M}_v(\beta_v) \times_L \partial C_2(L) \times_L \mathcal{M}_w(\beta_w)$ is

$$T\mathcal{M}_v \times_{TL} (\mathbb{R}_{\text{out}} \times T\partial C_2(L)) \times_{TL} T\mathcal{M}_w = \mathbb{R}_{\text{out}} \times T\mathcal{M}_v \times_{TL} T\partial C_2(L) \times_{TL} T\mathcal{M}_w$$

as oriented space.

Let $\tilde{\mathcal{M}}(C)$ be the space of holomorphic maps from the unit disk representing the homology class $C \in H_2(M, L)$. Then

$$\mathcal{M}_1 = (\tilde{\mathcal{M}} \times S^1)/\mathrm{PSL}(2,\mathbb{R})$$

where $\mathrm{PSL}(2,\mathbb{R})$ is the automorphism group of the disk. Let $\phi_i^v \in T(\mathrm{PSL}(2,\mathbb{R}))$ be the infinitesimal element that fixes $1, -1 \in D_v$ and move counter-clockwise $i \in D_v$ and let $\phi_{-1}^v \in T(\mathrm{PSL}(2,\mathbb{R}))$ be the infinitesimal element that fixes $i, 1 \in D_v$ and move counter-clockwise $-1 \in D_v$. As oriented spaces we have

$$T\tilde{\mathcal{M}}_v = T\mathcal{M}_v \times \{\phi_i^v, \phi_{-1}^v\}.$$

Let $\phi_{+1}^w, \phi_i^w \in \mathrm{PSL}(2,\mathbb{R})$ be the analogous elements for D_w. As oriented spaces we have

$$T\tilde{\mathcal{M}}_w = T\mathcal{M}_w \times \{\phi_{+1}^w, \phi_i^w\}.$$

The fiber product $\tilde{\mathcal{M}}_v \times_L \tilde{\mathcal{M}}_w$ is made using the evaluation map on $1 \in D_v$ and $-1 \in D_w$. The orientation in its tangent space is given by

$$\begin{aligned} T\tilde{\mathcal{M}}_v \times_{TL} T\tilde{\mathcal{M}}_w = \{\phi_i^v, \phi_{-1}^v\} \times T\mathcal{M}_v \times_{TL} T\mathcal{M}_w \times \{\phi_{+1}^w, \phi_i^w\} = \\ -\mathbb{R}_{\text{out}} \times T\mathcal{M}_v \times_{TL} T\mathcal{M}_w \times \{\phi_{-1}^v, \phi_{+1}^w, \mathbb{R}_{\text{tot}}\}. \end{aligned}$$

Here we have used that $\{\phi_i^v, \phi_i^w\} = \{\mathbb{R}_{\text{out}}, \mathbb{R}_{\text{tot}}\}$. Observe that $\{\phi_{-1}^v, \phi_{+1}^w, \mathbb{R}_{\text{tot}}\}$ is the orientation of $\mathrm{PSL}(2,\mathbb{R})$ acting on $\tilde{\mathcal{M}}_v \times_L \tilde{\mathcal{M}}_w$.

By Lemma 46.5 of [2], the gluing map $\tilde{\mathcal{M}}_v \times_L \tilde{\mathcal{M}}_w \to \tilde{\mathcal{M}}$ is orientation preserving in the sense of Kuranishi structures. The lemma follows.

□

3. Systems of singular chains

For each decorated tree T, let $C_T(L)$ be the orbifold

$$C_T(L) = \left(\prod_{e \in E(T)} C_e(L) \right) / \mathrm{Aut}(T).$$

The boundary of $C_T(L)$ can be decomposed in boundary faces corresponding to isomorphims classes of pairs (T, e)

$$\partial_e C_T(L) = \left(C_e(L) \times \prod_{e' \neq e} C_{e'}(L) \right) / \mathrm{Aut}(T, e)$$

where e is an internal edge of T. Observe that since $\partial C_2(L) \to L$ is an S^2-fibration, the induced map

$$\partial_e C_T(L) \to L \times C_{T/e}(L) \tag{12}$$

is an S^2-fibration.

Definition 3. *A system of singular chains*

$$W_{\mathcal{T}} = \{W_T\}_{T \in \mathcal{T}}$$

is a collection of singular chains $W_T \in C_{|E(T)|}(C_T(L), o_T)$ with twisted coefficients in o_T. We consider two singular chains equal if they are the same as currents. We assume the following properties.

(a) *For each $T \in \mathcal{T}$, W_T intersects transversely the boundary of $C_T(L)$. For each internal edge e define $\partial_e W_T = W_T \cap \partial_e C_T(L)$.*

(b) *There exist singular chains*

$$\partial'_e W_T \in C_{|E(T)|-3}(C_{T/e}(L), o_T)$$

such that $\partial_e W_T$ is the geometric preimage of $\partial'_e W_T$ over the S^2-fibration (12) (remember that the identities are considered identity of currents).

(c) *The following identity holds as currents*

$$\partial W_T = \sum_{v \in V(T)} \partial_v W_T + \sum_{e \in E(T)} \partial_e W_T. \tag{13}$$

Here $\partial^0_e W_T$ is the image of $\partial'_e W_T$ using the projection $L \times C_{T/e}(L) \to C_{T/e}(L)$ and

$$\partial_v W_T = \sum_{(T'/e, e/e) = (T, v)} \partial^0_e W_{T'}.$$

3.1. **Homotopies.** An homotopy $Y_{\mathcal{T}}$ between $W_{\mathcal{T}}$ and $W'_{\mathcal{T}}$ is a collection of singular chains

$$Y_T \in C_{|E(T)|+1}([0,1] \times C_T(L), o_T)$$

that satisfies analogous properties as Definition 3 with formula (13) replaced by

$$\partial Y_T = \sum_{v \in V(T)} \partial_v Y_T + \sum_{e \in E(T)} \partial_e Y_T + \{0\} \times W_T - \{1\} \times W'_T \tag{14}$$

Suppose that $Y_{\mathcal{T}}$ and $X_{\mathcal{T}}$ are two homotopies between $W_{\mathcal{T}}$ and $W'_{\mathcal{T}}$. We say that $Y_{\mathcal{T}}$ is equivalent to $X_{\mathcal{T}}$ if there exists a collection of singular chains $Z_{\mathcal{T}}$ with

$$Z_T \in C_{|E(T)|+2}([0,1]^2 \times C_T(L), o_T)$$

that satisfies analogous properties as Definition 3 with formula (13) replaced by

$$\partial Z_T = \sum_{v\in V(T)} \partial_v Z_T + \sum_{e\in E(T)} \partial_e Z_T + [0,1]\times\{0\}\times W_T - [0,1]\times\{1\}\times W'_T + \{0\}\times Y_T - \{1\}\times X_T \tag{15}$$

4. Invariance

4.1. **Perturbation of the Kuranishi structure.**

We want to use Lemma 1 in order to identify in pairs the boundaries of the $\overline{\mathcal{M}}_T$ for different T.

The only boundary face of $\partial\overline{\mathcal{M}}_{\mathcal{T}}$ with no cousin is the last set in the relation (10). Assume that the homological class of L in M is trivial. Let $B \in C_4(M)$ such that $\partial B = L$. For a fixed B we can consider the fiber product

$$\overline{\mathcal{M}}_{0,1} \times_M B. \tag{16}$$

The boundary of the Kuranishi space (16) can be identified with the last set in relation (10)

$$\partial(\overline{\mathcal{M}}_{0,1} \times_M B) \cong \overline{\mathcal{M}}_{0,1} \times_M L. \tag{17}$$

Denote by s_n a perturbation of the Kuranishi structure of $\overline{\mathcal{M}}_T$ for each decorated tree T and $\overline{\mathcal{M}}_{0,1} \times_M B$.

Definition 4. *We say that s_n is coherent in the boundary if for each tree T and for each vertex $v \in V(T)$, s_n respects the isomorphism of Lemma 1. This means that the pull-back of the restriction of s_n on $S(\partial_v\overline{\mathcal{M}}_T)$ using the map (9) coincides with the restriction of s_n to $\partial_e\overline{\mathcal{M}}_{T'}$. In the case of the tree T_0 without edges we need to impose also that the perturbation on $\partial_v\overline{\mathcal{M}}_T$ and $\overline{\mathcal{M}}_{0,1} \times_M B$ agree on $\overline{\mathcal{M}}_{0,1} \times_M L$.*

Fix a perturbation s_n of the Kuranishi structure that satisfies condition (4). The existence of such a perturbation can be proved using the standard machinery developed in Section 6 of [1] or Appendix A of [2] (see Lemma 5 below).

There exists a natural strongly continuous map

$$\mathrm{ev} : \overline{\mathcal{M}}_T(J) \to C_T(L). \tag{18}$$

As in formula (6.10) of [1], the strongly continuous map (18) defines a singular chain

$$W_T = \mathrm{ev}_*((s_n|^{-1}_{\partial\overline{\mathcal{M}}_T})(0)) \tag{19}$$

that is a chain in $C_T(L)$ with twisted coefficients in o_T.

If $T = T_0$ is the tree with one vertex and no edges we need to add a correction to formula (19):

$$W_{T_0} = \mathrm{ev}_*((s_n|^{-1}_{\partial\overline{\mathcal{M}}_{T_0}})(0)) + \mathrm{ev}_*((s_n|^{-1}_{\overline{\mathcal{M}}_{0,1}\times_M B})(0)) \in \mathbb{Q}. \tag{20}$$

The target of the map ev in formula (20) is a point.

4.2. **Invariance.**

The Kuranishi structure we constructed may depend on the various choices we made. However we have the following (the proof is minor adaptation of the proof of Theorem 17.11 of [1]):

Proposition 4. *The system of singular chains $W_{\mathcal{T}}$ depends on the almost complex structure J and various choice we made to define a Kuranishi structure. Different choices lead to system of homological chains that are homotopic, with the homotopy determined up to equivalence.*

Proof. Let J and J' be two different complex structures compatible with the symplectic structure ω. Let J_s be a family of compatible almost complex structures such that $J_s = J$ for $s \in [0, \varepsilon]$ and $J_s = J'$ for $s \in [1-\varepsilon, 1]$. Define

$$\overline{\mathcal{M}}_{\mathcal{T}}(J_{\text{para}}) = \cup_{s\in[0,1]}\{s\} \times \overline{\mathcal{M}}_{\mathcal{T}}(J_s).$$

As in Theorem 17.11 of [1] we can endow $\overline{\mathcal{M}}_{\mathcal{T}}(J_{\text{para}})$ with a compact and Hausdorff topology. Moreover there exists a Kuranishi structure on $\overline{\mathcal{M}}_{\mathcal{T}}(J_{\text{para}})$ that extends the Kuranishi structure of $\overline{\mathcal{M}}_T(J)$ and $\overline{\mathcal{M}}_T(J')$.

Lemma 5 implies that there exist a transverse multisections $s_{p,n}$ perturbing s_p and satisfying the coherence condition of Definition 4. Moreover if we start with a perturbation of $\overline{\mathcal{M}}_{\mathcal{T}}(J)$ and $\overline{\mathcal{M}}_{\mathcal{T}}(J')$, $s_{p,n}$ can be constructed such that it extends these perturbations.

For each tree T, define the chain $\tilde{W}_T$ in $C_T(L) \times [0, 1]$ as in (19) and (20). Lemma 4.7 of [1] implies that $\partial\tilde{W}_T$ is the intersection with the boundary of $\overline{\mathcal{M}}_T(J_{\text{para}})$:

$$\partial\tilde{W}_T = \text{ev}_*((s_n|^{-1}_{\partial\overline{\mathcal{M}}_T(J_{\text{para}})})(0))$$

The system of chains $\tilde{W}_{\mathcal{T}} = \{\tilde{W}_T\}$ defines an homotopy between the system of chains $W_{\mathcal{T}}$ and $W'_{\mathcal{T}}$.

□

Lemma 5. *There exists a perturbation s_n of the Kuranishi structure of $\overline{\mathcal{M}}_T(J)$ coherent in boundary in the sense of Definition 4 and transverse to the zero section.*

Proof. The construction of the multi sections $s_{p,n}$ is done using the standard machinery of Kuranishi structures developed in [1] or Appendix A of [2]. The construction is done by induction on the number of edges of the tree. In each step we impose the conditions of Definition 4.

The Kuranishi structure of $\overline{\mathcal{M}}_T(J_{\text{para}})$ is defined by a set of Kuranishi charts $(V_p, E_p, \Gamma_p, s_p, \psi_p)$. A minor adaptation of the proof of Lemma 6.3 of [1] gives a good cover extending a good covering of $\partial\overline{\mathcal{M}}_T(J_{\text{para}})$.

Suppose first that T is just a vertex. In this case $\overline{\mathcal{M}}_T$ is the usual space of disks, therefore it is standard to construct a multi-section s_n transverse to the zero section.

Now consider a tree T with at least one internal edge and assume that the multi-section s_n has been constructed for all the trees with less edges than T. Let $e \in E(T)$. The condition of Definition 4 (a) define a multisection on $\partial_e\overline{\mathcal{M}}_T$ in terms of the multi-section of $\partial_v\overline{\mathcal{M}}_{T/e}$. These sections are compatible in the corners. This follow by the induction hypothesis. If $e_1, e_2 \in E(T)$ then the section on $\partial_{e_1}\overline{\mathcal{M}}_{\ T}$ restricted to $\partial_{e_1,e_2}\overline{\mathcal{M}}_T$ is constructed from the section on $\partial_{v_1,v_2}\overline{\mathcal{M}}_{T/\{e_1,e_2\}}$. This is the same for the section on $\partial_{e_2}\overline{\mathcal{M}}_T$ restricted to $\partial_{e_1,e_2}\overline{\mathcal{M}}_T$.

For each p, this multi-section gives a multi-section defined on the union of some closed boundary face of V_p. Since V_p is a manifold with corners, it is not hard to extend this multi-section to a neighborhood of these faces. Lemma 17.4 of [1] or Theorem A1.23 of [2] can be used to give a transverse multi-section of $\overline{\mathcal{M}}_T(J, \beta)$. □

5. Linking number and Gromov-Witten invariants

In this section we assume that each connected component of L has the rational homology of a sphere. Let $\mathcal{T}_0$ be the set decorated trees with no external edges .

We want to extend the definition of linking number of two curves to system of chains. Remember that to define the linking of two curves γ_1 and γ_2 in L we can proceed as follows. The product $\gamma_1 \times \gamma_2$ defines a 2-chain in $L \times L$ homological trivial. Let C be a 3-chain in $L \times L$ such that $\partial C = \gamma_1 \times \gamma_2$ and is transversal to diagonal. The linking number of γ_1 and γ_2 is given by the intersection number of C with the diagonal. This argument can be straightforwardly extended to systems of chains.

Lemma 6. *There exists a system of chains $W'_{\mathcal{T}_0}$ that is homotopic to $W_{\mathcal{T}_0}$ such that*

$$W'_T = 0$$

if T has at least an edge. Moreover $W'_{\mathcal{T}_0}$ is unique.

Proof. It can be useful to observe that systems of chains can be equivalently defined as singular chains in $\{L^{H(T)}/\mathrm{Aut}(T)\}_T$ that are transverse to the diagonals associated with the edges of the trees and moreover satisfy analogous condition of coherence formulated in terms of the intersections with the diagonals.

Let $T_1 \in \mathcal{T}_0(A)$ be the decorated tree in the homological class A with maximal number of edges. We can construct an homotopy between $W_{\mathcal{T}_0(A)}$ and a system of chains with $W_{T_1} = 0$ as follows. W_{T_1} is a singular chain in $L^{H(T_1)}/\mathrm{Aut}(T_1)$ homological trivial. This can be used to construct an homotopy between W_T and zero chain and it can be extended to all the trees using the compatibility condition in the boundary. This yields to an homotopy between $W_{\mathcal{T}}$ and a system of chains with T_1-component zero. We can proceed recursively and construct the required homotopy.

Suppose now that $W''_{\mathcal{T}_0}$ is another system of chains as in the Lemma. There exists an homotopy $Z_{\mathcal{T}_0}$ between $W'_{\mathcal{T}_0}$ and $W''_{\mathcal{T}_0}$. By the same argument above $Z_{\mathcal{T}_0}$ is equivalent to an homotopy with $Z_T = 0$ for all the trees T with at least one edge. From this it follows that $W'_{\mathcal{T}_0} = W''_{\mathcal{T}_0}$. □

The last lemma allows us to define the linking number of $W_{\mathcal{T}_0(A)}$ by

$$\mathrm{link}(W_{\mathcal{T}_0(A)}) = W'_{T_0} \in \mathbb{Q}.$$

Here T_0 is the trivial tree with no edges and exactly one vertex, so that $C_{T_0}(L)$ is a point.

Theorem 7. *The rational number link$(W_{\mathcal{T}_0(A)})$ does not depend on the almost complex structure and various choice we made to define a Kuranishi structure.*

Proof. The theorem follows directly from Proposition (4). □

We can now define the Open Gromov-Witten invariants that counts multi-disks in the relative homology class $A \in H_2(X, L)$ as

$$F_{0,1}(A) = \mathrm{link}(W_{\mathcal{T}_0(A)})$$

It also easy to see how $F_{0,1}(A)$ depends on the singular chain B that we have used to define (16). Assume that $B' \in C_4(M)$ is another singular chain with $\partial B' = L$. Since $B - B'$ is a cycle we have

$$F_{0,1}(A) - F_{0,1}(A)' = \mathrm{ev}_*([\overline{\mathcal{M}}_{0,1}(A)]) \cap (B - B'). \tag{21}$$

6. Higher genus

In this section we assume that L is spin so that all the moduli spaces we consider are oriented (see [5]).

6.1. **Multi-curves.** Let $\mathcal{M}_{(g,h),(n,\overrightarrow{m})}(A)$ be the Kuranishi space of stable maps of type (g,h) with n internal marked points and $\overrightarrow{m}=(m_1,...,m_h)$ boundary marked points, representing the relative homology class $A \in H_2(M,L)$. This space has been studied in [5].

Definition 5. *A decorated graph G is a ribbon graph endowed with the following data*

- *A set $I(G)$. For each $i \in I(G)$ a relative homological classes $A_i \in H_2(M,L)$ and an oriented surface Σ_i . Let g_i be the genus, h_i the number of boundary components, n_i be the number of internal marked points, $\overrightarrow{m_i}$ be the number of boundary marked points of Σ_i. If $A_i=0$ we assume that Σ_i is stable.*
- *A one to one correspondence between vertices of G and the boundary components of $\{\Sigma_i\}_{i\in I}$.*
- *For each $v \in V(G)$, a one to one correspondence between the half edges $H(v)$ starting in v and the boundary marked points of the boundary component associated to v.*

The homology class of G is given by $A=\sum_{i\in I} A_i$. From a decorated graph G we can construct a surface Σ_G replacing each edge of G with a strip respecting the orientation of the surfaces Σ_i. We say that the decorated graph G is connected if the surfaces Σ_G is connected.

Denote by $\mathcal{G}_{(g,h),(n,\overrightarrow{m})}(A)$ the set of connected decorated graphs G where Σ_G has genus g, h boundary components, n internal marked points, $\overrightarrow{m}$ external edges, representing the relative homology class $A \in H_2(M,L)$.

For each $i \in I$ denote $\overline{\mathcal{M}}_i = \overline{\mathcal{M}}_{(g_i,h_i),(n_i,\overrightarrow{m}_i)}(A_i)$. The evaluation map on the punctures defines a map

$$\prod_{i\in I(G)} \overline{\mathcal{M}}_i \to L^{H(G)} \times M^n. \tag{22}$$

We have also the map

$$\left(\prod_{e\in E(T)} C_e(L)\right) \times B^n \to L^{H(G)} \times M^n. \tag{23}$$

The moduli space of multi-curves $\overline{\mathcal{M}}_G$ is defined as the fiber product of the maps (22) and (23) modulo the automorphism group of the decorated graph G:

$$\overline{\mathcal{M}}_G = \left(\left(\prod_{i\in I}\overline{\mathcal{M}}_i\right) \times_{(L^{H(G)}\times M^n)} \left(\prod_{e\in E(G)} C_e(L)\times B^n\right)\right)/\mathrm{Aut}(G).$$

6.2. **Boundary.** In [5] is proved that the boundary $\overline{\mathcal{M}}_{(g,h),(n,\overrightarrow{m})}$ can be subdivided in components of type E, $H1$, $H2$, $H3$. Each boundary face is associated to a boundary component of the surface. For each boundary component C, denote by

$\partial_C\overline{\mathcal{M}}_{(g,h),(n,\vec{m})}$ the union of the boundary faces of $\overline{\mathcal{M}}_{(g,h),(n,\vec{m})}$ associated to a degeneration of C. We then have

$$\partial\overline{\mathcal{M}}_{(g,h),(n,\vec{m})} = \bigsqcup_C \partial_C\overline{\mathcal{M}}_{(g,h),(n,\vec{m})}.$$

Let G be a decorated graph. For each $v \in V(G)$ define

(24)
$$\partial_v\overline{\mathcal{M}}_G = \left(\left(\partial_v\overline{\mathcal{M}}_i \times \prod_{i'\neq i}\overline{\mathcal{M}}_{i'}\right) \times_{(L^{H(G)}\times M^n)} \left(\prod_e C_e(L) \times B^n\right)\right)/\mathrm{Aut}(G,v)$$

where $i \in I(G)$ is the surface corresponding to v. For each $e \in E(G)$ define

(25)
$$\partial_e\overline{\mathcal{M}}_G = \left(\left(\prod_{i\in I(G)}\overline{\mathcal{M}}_i\right) \times_{\left(L^{H(G)}\times M^n\right)} \left(\partial C_e(L) \times \prod_{e'\neq e} C_{e'}(L) \times B^n\right)\right)/\mathrm{Aut}(G,e).$$

Finally for each internal marked point p define

(26)
$$\partial_p\overline{\mathcal{M}}_G = \left(\left(\prod_{i\in I}\overline{\mathcal{M}}_i\right) \times_{(L^{H(G)}\times M^n)} \left(\prod_{e\in E(G)} C_e(L) \times (B^{n-1}\times L)\right)\right)/\mathrm{Aut}(G,p)$$

The Kuranishi spaces (24), (25) and (26) are the boundary faces of $\overline{\mathcal{M}}_G$.

Definition 6. *Let $e \in E(G)$ be an internal edge of G. Let v, w be the vertices of e that are boundary components of $i_v, i_w \in I(G)$ respectively. The decorated graph G/e is defined by contracting the edge e. The surfaces Σ_{i_v} and Σ_{i_w} are replaced by the surface made by gluing the boundary marked points associated to e and smoothing the resulting node.*

Observe that in the preview definition we can have $i_v = i_w$ or even $v = w$. In this process we get a boundary node of type

- H1 if $v = w$,
- H2 if $v \neq w$ but $i_v = i_w$,
- H3 if $i_v \neq i_w$.

The boundary nodes of type E are considered in the following:

Definition 7. *Let $v \in V(G)$ be a vertex of G corresponding to a boundary component of Σ_i without marked points. The decorated graph G/v is the graph G with the vertex v removed and with the correspondent boundary component of Σ_i replaced with an internal marked point.*

With this notation we can formulate the generalization of Lemma 1:

Lemma 8. *Let G be a decorated graph. For each $v \in V(G)$ we have*

$$S(\partial_v\overline{\mathcal{M}}_G) \cong \Big(\bigsqcup_{(G'/e,e/e)=(G,v)} \partial_e\overline{\mathcal{M}}_{G'}\Big) \sqcup S(\partial_{v/v}\overline{\mathcal{M}}_{G/v}). \tag{27}$$

The last term is present only if v is an isolated vertex. v/v denotes the internal marked point of G/v that we get from v.

6.3. **Systems of chains.** As in the genus zero we have a natural map

$$\mathrm{ev} : \overline{\mathcal{M}}_G \to C_G(L)$$

where $C_G(L)$ is the orbifold

$$C_G(L) = \left(\prod_{e \in E(G)} C_e(L) \right) / \mathrm{Aut}(G). \tag{28}$$

Using the isomorphism (27) it is easy to extend the definition of coherent perturbation in the boundary to $\overline{\mathcal{M}}_{\mathcal{G}}$. Each such perturbation induces a system of chains

$$W^0_{\mathcal{G}} = \{W^0_G\}_{G \in \mathcal{G}}$$

where for each G W_G is a singular chain of $C_G(L)$.

Let $\mathcal{G}_0$ be the set of decorated graphs without internal marked points and without external edges. Define

$$W_G = \sum_{G'} W^0_{G'}$$

where the sum is over all the decorated graphs G' made from G substituting with internal marked points some boundary component of G with no marked points (observe that the sum makes sense since $C_{G'}(L) \cong C_G(L)$ for all such graphs G'). We therefore have a system of chains

$$W_{\mathcal{G}_0} = \{W_G\}_{G \in \mathcal{G}_0}.$$

The definition of homotopy and equivalence we gave for decorated trees extended naturally to decorated graphs.

Lemma 9. *Different choices of coherent pertubations lead to homotopic systems of chains, with homotopy unique up to equivalence.*

6.4. **Gromov-Witten invariants.** Assume that each connected component of L has the rational homology of a sphere. As in the case of disks we have:

Lemma 10. *There exists a system of chains $W'_{\mathcal{G}_0}$ that is homotopic to $W_{\mathcal{G}_0}$ such that*

$$W'_G = 0$$

for all $G \in \mathcal{G}_0$ with at least one edge. Moreover $W'_{\mathcal{G}_0}$ is unique.

For each (g, h) let $G^0_{g,h}(A) \in \mathcal{G}_0$ be the unique decorated graph with no edges of genus g with h boundary components and in relative homology class $A \in H_2(X, L)$ ($G^0_{g,h}(A)$ represents an ordinary surface). By definition $C_{G_0^{g,h}}(L)$ is given by a point. Define the Gromov-Witten invariant as the rational number

$$F_{g,h}(A) = W'_{G^0_{g,h}(A)} \in \mathbb{Q}.$$

From Lemma 9 and Lemma 10 follows:

Theorem 11. *The rational numbers $F_{g,h}(A)$ do not dependent on the almost complex structure and various choices we made to define the Kuranishi structure.*

Observe that the rational numbers $F_{g,h}(A)$ depend not only on the pair (X, L) by also on the choice of the singular chain B. However, for different choices of B the invariants can be expressed by a formula analogous to (21).

REFERENCES

[1] K. Fukaya; K. Ono, *Arnold conjecture and Gromov-Witten invariant*, Topology 38 (1999), no. 5, 933–1048.

[2] K. Fukaya, Y.-G. OH, H. Ohta, K. Ono, *Lagrangian intersection Floer theory - anomaly and obstruction*.

[3] V. Iacovino, Open Gromov-Witten theory on Calabi-Yau three-folds II, arXiv:0908.0393.

[4] V. Iacovino, Frame ambiguity in Open Gromov-Witten invariants, preprint.

[5] M. Liu, *Moduli of J-Holomorphic Curves with Lagrangian Boundary Conditions and Open Gromov-Witten Invariants for an* S^1*-Equivariant Pair*, arXiv:math/0210257.

[6] J. P. Solomon *Intersection theory on the moduli space of holomorphic curves with Lagrangian boundary conditions*, arXiv:math/0606429.

[7] E. Witten, *Chern-Simons Gauge Theory As A String Theory*, arXiv:hep-th/9207094.

MAX-PLANCK-INSTITUT FÜR MATHEMATIK, BONN
E-mail address: `iacovino@mpim-bonn.mpg.de`